

C  
540  
WE.E

**ANALYSIS OF CADMIUM IN DRINKING WATER  
IN PADAVI SIRIPURA AREA  
AND  
MODIFICATION OF DIPHENYLTHIOCARBAZONE**

**A PROJECT REPORT PRESENTED BY**

**DONA PABASARA PRIYANGANI WEERASINGHE**

to the Board of Study in Chemical Sciences of the  
**POSTGRADUATE INSTITUTE OF SCIENCE**

*in partial fulfillment of the requirement  
for the award of the degree of*

**MASTER OF SCIENCE IN ANALYTICAL CHEMISTRY**

of the

**UNIVERSITY OF PERADENIYA**


**SRI LANKA**

**2004**

**580454**

**ANALYSIS OF CADMIUM IN DRINKING WATER  
IN PADAVI SRIPURA AREA  
AND  
MODIFICATION OF DIPHENYLTHIOCARBAZONE**

**D.P.P. Weerasinghe**  
Department of Chemistry  
University of Peradeniya  
Peradeniya  
Sri Lanka



A significant increase in the number of patients with Chronic Renal Failure (CRF) in Padavi Sripura area in the North Central Province (NCP) of Sri Lanka has been reported. It was suspected that this was due to drinking of cadmium contaminated water because the symptoms were similar to that of cadmium poisoning. Analysis of water samples collected from Padavi Sripura area was carried out with a view to testing this proposition. An analysis using a Graphite Furnace Atomic Absorption Spectrometer (GFAAS) did not reveal the presence of cadmium in concentrations sufficient to pose a health hazard. However the effects of bioaccumulation have to be carefully investigated before a final conclusion is reached.

An attempt was made to develop a complexing agent for the determination of cadmium with improved sensitivity by chemically modifying diphenylthiocarbazon. Bamberger method and the Fischers method used for the synthesis of diphenylthiocarbazon were modified and used in order to synthesis di-(2,4-dinitrophenyl)thiocarbazon and di-(4-nitrophenyl)thiocabazon. However this part of the project could not be completed due to various constraints such as the nonavailability of the appropriate starting materials. The coupling reaction of the Bamberger method occurred smoothly to give 1,5-(4-nitrophenyl)nitroformazan in good yield but the conversion of 1,5-(4-nitrophenyl)nitroformazan to di-4-nitrophenylthiocarbazon with reducing agent, ammonium hydrogensulphide was found to be ineffective. First step of the modified Fischers method gave a product only in very small quantities. This may be due to the deactivating effect of the nitro functionality.