



**Investigation on the Involvement of Bacteria in  
Diseases of Cultured Shrimp in Sri Lanka**

Panawala Vidanelage Sachie Panawala


For the **Degree of Master of Philosophy**

**University of Peradeniya**

Department of Veterinary Pathobiology

Faculty of Veterinary Medicine and Animal Science

**August – 2007**



**619288**

## Abstract

Shrimp culture, in spite of its relatively short history as a commercial industry is developing at a rapid pace, but the sustainability of this trade is now at stake as threats of disease outbreaks have become a great concern of the shrimp producing countries including Sri Lanka. Various kinds of bacterial species act as causative agents in many of these diseases. Many other countries have studied these pathogenic bacteria in relation to diseases of shrimp but information on Sri Lankan scenario is not available. Therefore, a study was designed with the intension of addressing this problem. As background knowledge on farm management is a prerequisite to study diseases and their causative agents, information on randomly selected farms along the Western coastal belt of Sri Lanka was gathered. Most of the farms were found to be following malpractices in view of health considerations. Non-availability of sedimentation tanks for effluents and direct release of wastewater to water sources are among them. The water quality parameters of the ponds and water sources that supply them were generally within the recommended levels except for the total suspended solids. The mean ammonia concentration in pond water was remarkably higher than the standard levels. The mean dissolved oxygen levels of pond-water and water sources were at the lower limit of recommended range and this can be improved by increasing the frequency of aeration. Haemolymph from diseased shrimp and water samples were subjected to bacteriological investigations. As a result, five *Vibrio* species namely, *Vibrio harveyi*, *V. parahaemolyticus*, *V. vulnificus*, *V. alginolyticus* and *V. anguillarum*, along with *Staphylococcus aureus* were isolated and identified. Conventional bacteriological methods were employed to identify them. Immunohistochemistry, which is a more sensitive and confirmative method, confirmed identification of these bacteria and along with histological studies, provided evidence of their active

involvement in diseases of shrimp. Normal healthy shrimp were challenged with the isolated bacteria at a dose of  $2 \times 10^7$  CFU per shrimp and all of them showed significant ( $p < 0.05$ ) mortality with *Vibrio alginolyticus* being the most virulent. There were no deaths after first 3 days of the challenge study. Antimicrobial susceptibility of the isolates was tested and all showed resistance to erythromycin. Emerging resistance to tetracycline was also clearly evident. The efficacy of the commonly used remedial measures was tested by an *in vitro* study, and the use of probiotics and exposure of water to sunlight proved to be useful, while applying sugar provided evidence to be not effective in controlling the bacterial growth. Identifying probiotics, formulating quality feed, identifying proper systems for effluent management, development of successful vaccines and selection and development of disease tolerant shrimp are the areas that warrant future research.

	Viral diseases	18
	White Spot Syndrome (WSS)	19
	Yellow Head Disease (YHD)	19
	Bacterial diseases	20
	Vibriosis	21
	Poor water quality	22
	Effects on the environment	23
	Use of antimicrobials	23
	Investigation on the involvement of bacteria in diseases of cultured shrimp in Sri Lanka	24
1	Shrimp Farming Practices in the North Western Coastal Belt of Sri Lanka	
	Introduction	26
	Materials and Methods	27
	Results	28
	Discussion	33
	Conclusion	39
2	A Study on Water Quality of Selected Shrimp Farms and Main Water Sources that Supply Them	
	Introduction	41
	Materials and Methods	43
	Results	44
	Discussion	44
	Conclusion	49