EFFECT OF MANNAN OLIGOSACCHARIDES ON SOME NUTRITIONAL AND BIOCHEMICAL PARAMETERS OF MICE

i

A PROJECT REPORT PRESENTED BY

THARAKA ROSHANI KUMBUKAGE

to the Board of Study in Biochemistry and Molecular Biology
POST GRADUATE INSTITUTE OF SCIENCE

in partial fulfillment of the requirement for the award of the degree of

MASTER OF SCIENCE IN EXPERIMENTAL BIOTECHNOLOGY

of the

UNIVERSITY OF PERADENIYA SRI LANKA 2009



EFFECT OF MANNAN OLIGOSACCHARIDES ON SOME NUTRITIONAL AND BIOCHEMICAL PARAMETERS OF MICE

T.R. Kumbukage

Department of Biochemistry,
Faculty of Medicine,
University of Peradeniya,
Peradeniya.

Prebiotics are non-digestible food ingredients that beneficially affect the host by selectively stimulating the growth and/or activity of one or a limited number of bacteria in the colon, and thus improve host health. They have become increasingly popular as feed additives in recent years due to their numerous health benefits. Mannanoligosaccharides, a prebiotic candidate derived from the cell wall of Saccharomyces cerevisiae have shown promising effects on animal health such as decreasing pathogenic microflora in the gut, stimulating a strong immune response and elevating the strength of intestinal mucosa. The aim of this study was to determine the effect of addition of mannan-oligosachcharide Bio-Mos®, on some nutritional and biochemical parameters of mice. Thirty-two, female, sixteen to twenty-week-old, Balb/c mice were assigned randomly into two groups, namely treatment and control, with eight replicates of 4 mice each. Both groups were fed with broiler starter ration ad libitum, but to the feed of the treatment group 5% of Bio-Mos® was added. Body weights were recorded throughout the experimental period. Blood was collected from mice after a period of one-month feeding schedule. Serum biochemical parameters including total cholesterol, HDL-cholesterol, total protein and albumin levels were determined using diagnostic kits. A differential count of the white blood cells was also performed. Bio-Mos® supplementation had significantly reduced the total serum cholesterol level of test group mice. Both HDL cholesterol level and cholesterol/HDL ratio were not significantly affected by Bio-Mos® supplementation. Although the total serum protein concentration of mice was not significantly affected by Bio-Mos® supplementation, serum albumin level had shown a significant increase (P < 0.05). An improvement of the body weight gain was observed in test group mice, but with no statistically significant difference. Addition of Bio-Mos®

favourably increased the lymphocyte count and significantly decreased in the neutrophil count (P<0.05) of mice. It was concluded that inclusion of prebiotic (Bio-Mos®) to the feed of mice had a positive influence on some biochemical and immunological indices of blood and seems to be a suitable functional feed additive from the production and animal health point of view.

